**Archive of Applied Mechanics manuscript No.**
(will be inserted by the editor)

**Steffen Strauch for the CLAS Collaboration**

# Baryon spectroscopy with polarization observables from CLAS

arXiv:1602.07284v1 [nucl-ex] 23 Feb 2016

Received: date / Accepted: date

**Abstract** The spectrum of nucleon excitations is dominated by broad and overlapping resonances. Polarization observables in photoproduction reactions are key in the study of these excitations. They give indispensable constraints to partial-wave analyses and help clarify the spectrum. A series of polarized photoproduction experiments have been performed at the Thomas Jefferson National Accelerator Facility with the CEBAF Large Acceptance Spectrometer (CLAS). These measurements include data with linearly and circularly polarized tagged-photon beams, longitudinally and transversely polarized proton and deuterium targets, and recoil polarizations through the observation of the weak decay of hyperons. An overview of these studies and recent results will be given.

**Keywords** Baryon Resonances · Spectroscopy · Polarization Observables

## 1 Introduction

The nucleon is a color-neutral object which consists of color-charged quarks and gluons. Quantum chromodynamics (QCD) is the fundamental theory of the strong interaction between quarks and gluons. Valuable information about QCD can be learned from nuclear spectroscopy; *e.g.* information about the internal degrees of freedom in a nucleon. Experiments, especially pion-nucleon scattering, have confirmed low-lying excited states predicted by quark models with three independent quark degrees of freedom. These models, however, predict an overabundance of higher-lying excited states compared to what has been observed until now [1]. Also, recent lattice QCD calculations [2] find a large number of not-yet-discovered nucleon resonances. To clarify the nucleon resonance spectrum is an important task in the study of QCD.

Nucleon resonances are short-lived, and the identification of resonances in partial-wave analyses of experimental data is complicated by their large width and overlap. Experimental cross sections are insufficient, and polarization observables are crucial to constrain these analyses. A complete set of certain polarization observables is necessary to unambiguously determine the amplitudes of the reaction. In the photoproduction of pseudoscalar mesons, a formally complete experiment requires at each energy and angle the measurement of at least eight carefully chosen observables [3]. In the photoproduction of two mesons, even more observables are needed [4]. It is also important to include in the analysis data from a variety of excitation and decay channels, as some of the missing states may couple only weakly to, *e.g.*, the $\pi N$ channel.

Supported in parts by the U.S. National Science Foundation: NSF PHY-1505615.

S. Strauch
University of South Carolina
Tel.: +1-803-777-8197
Fax: +1-803-777-3065
E-mail: strauch@sc.edu

In the following, examples of recent photoproduction measurements of polarization observables from the CLAS Collaboration are presented. A similar overview has been given in [5]. These measurements include single pseudoscalar-meson, vector-meson, double-pion, and hyperon photoproduction off the proton and quasi-free off the bound neutron. The observables include single- and double-polarization observables with combinations of polarized beam, target, and the polarization of the recoiling baryon.

The experiments were performed in Experimental Hall B at the Thomas Jefferson National Accelerator Facility (JLab). The incident bremsstrahlung photon beams were energy-tagged [6] and either unpolarized, circularly, or linearly polarized. The photon beam irradiated the production target. Unpolarized liquid hydrogen and deuterium targets, as well as the newly developed polarized frozen-spin (FROST) [7] and HDice targets [8; 9], have been used in these experiments. Final-state particles were detected in the CEBAF Large Acceptance Spectrometer (CLAS) [10]. Recoil polarization was accessable in hyperon-production measurements through the measurement of the decay-proton angular distribution in the parity-violating weak decay of hyperons.

## 2 Unpolarized Targets

The CLAS Collaboration has studied the photoproduction reactions $\gamma p \to p\pi^0$ and $\gamma p \to n\pi^+$ with linearly-polarized photons in an energy range from 1.10 to 1.86 GeV [11]. The beam asymmetry observable $\Sigma$ has been obtained from the pion angular distributions with respect to the polarization direction of the linearly-polarized photon beam. The high statistics in these reactions allowed for precise constraints of partial-wave analyses. Resonance couplings have been extracted in fits of the SAID [12] analysis after including the new data set. The largest change from previous fits was found to occur for the 'well known' $\Delta(1700)3/2^-$ and $\Delta(1905)5/2^+$ resonances [11].

Polarization observables were also obtained in the hyperon-photoproduction reactions $\gamma p \to K^+\Lambda$ and $\gamma p \to K^+\Sigma^0$. Results include the $\Lambda$ recoil polarization $P$ from measurements with unpolarized photon beams [13; 14] as well as the polarization-transfer observables $C_x$ and $C_z$ with circularly-polarized beams [15]. These data were critical in a coupled-channel analysis by the Bonn-Gatchina group [16; 17]. In particular, the analysis found further evidence for the, at the time, poorly known $N(1900)3/2^+$ resonance. This resonance is predicted by symmetric three-quark models, but is not expected to exist in earlier quark-diquark models. Preliminary hyperon-photoproduction data with linearly-polarized photons off an unpolarized proton target have been obtained up to $W \approx 2.2$ GeV. Together with the recoil polarization of the hyperon, this gives access to five polarization observables: $\Sigma$, $P$, $T$, $O_x$, and $O_z$. Energy distributions of the preliminary results of the beam-recoil polarization observable $O_x$ for the $K^+\Lambda$ channel are shown in Fig. 1.

## 3 Polarized Targets

An integral part of the experimental N* program at JLab are experiments off polarized protons [18; 19; 20; 21; 22]. The Jefferson Lab frozen spin target (FROST) was constructed for use inside CLAS to allow for the measurement of polarization observables with longitudinally- or transversely-polarized protons in the butanol target material [7].

First results from the FROST program have been published for the $\gamma p \to \pi^+ n$ reaction [23]. The double-polarization observable $E$ has been determined in the energy range from 0.35 to 2.37 GeV from data of circularly-polarized photons incident on longitudinally-polarized protons. A subset of the about 900 data points is shown in the three energy bins of Fig. 2. Results from previous partial-wave analyses describe the new data at low photon energies reasonably well; at higher energies, however, significant deviations are observed. The data have been included in new analyses resulting in good descriptions of the data and in updated nucleon resonance parameters. One particularly interesting result is strengthened evidence for the poorly known $\Delta(2200)7/2^-$ resonance in improving the Bonn-Gatchina fit at the highest energies [24]. The mass of the $\Delta(2200)7/2^-$ resonance is significantly higher than the mass of its parity partner $\Delta(1950)7/2^+$, which is the lowest-mass $\Delta^*$ resonance with spin-parity $J^P = 7/2^+$.

The polarization observable $E$ has also been measured in the $\gamma p \to \eta p$ reaction from threshold to $W = 2.15$ GeV [27]. Because $\eta$ mesons have isospin zero, the reaction selects isospin-1/2 resonances


γ + p → K+ Λ
Observable: Beam-Recoil, Ox
-0.75 < cos θK < -0.55
-0.55 < cos θK < -0.35
-0.35 < cos θK < -0.15
-0.15 < cos θK < 0.05
0.05 < cos θK < 0.25
0.25 < cos θK < 0.45
0.45 < cos θK < 0.65
0.65 < cos θK < 0.85
W (GeV)

**Fig. 1** Preliminary energy distributions of CLAS $\gamma p \to K^+\Lambda$ data for the double-polarization observable $O_x$ in various kaon angular bins. Figure from D. Ireland (University of Glasgow).




**Fig. 2** CLAS data for the beam-target polarization observable $E$ in the $\gamma p \to \pi^+ n$ reaction for selected center-of-mass energy bins. The shaded bands indicate systematic uncertainties. The dashed curves in the upper panels are results from the SAID ST14 [12], Jülich14 [25], and BnGa11E [26] analyses. The solid curves in the lower panels are results from updated analyses including the new CLAS $E$ data. Figure from Ref. [23].

in the nucleon resonance spectrum. Figure 3 shows the data and a fit with the Jülich-Bonn dynamical coupled-channel model. It has been shown in Ref. [27] that the observable $E$ in $\eta$ photoproduction is especially suited to disentangle electromagnetic resonance properties. Initial investigation of these results show pronounced changes in the description of this observable when these new CLAS data are included. The fit in Fig. 3 describes the data quite well without the need for an additional narrow resonance near 1.68 GeV, which was previously suggested [27].




**Fig. 3** Observable $E$ in the $\gamma p \to \eta p$ reaction as a function of $W$ for various angular bins. The red curve shows fits with the Jülich-Bonn dynamical coupled-channel model. Figure from Ref. [27].

FROST data from other single-pion photoproduction channels and for other observables are under ongoing analyses. Examples of preliminary angular distributions for the target $T$ and beam-target $F$ polarization observables in the reaction $\gamma p \to \pi^0 p$ are shown in Fig. 4. These observables are extracted from data with unpolarized and circularly-polarized photons off transversally-polarized protons, respectively. In the figure, the data are compared with present results of partial-wave analyses which do not include these preliminary data.




**Fig. 4** Preliminary CLAS results for the observables $T$ and $F$ in the $\gamma p \to \pi^0 p$ reaction as a function of $\cos\theta_\pi$ for $W = 1.835$ GeV. The data are compared with various partial-wave analyses (not fitted to the data) from the SAID [12], MAID [28], and Bonn-Gatchina [26] groups. Figures from H. Jiang (University of South Carolina).

Hyperon photoproduction reactions are also studied with data from FROST. Figure 5 shows preliminary results of the beam-target observable $F$. The data cover energies between $W = 1.7$ GeV and $W = 2.3$ GeV. The data are compared to prior results of analyses with the RPR-Ghent [29], KAON-MAID [30], and Bonn-Gatchina [26] models. As none of the models describe the data well the FROST data will provide important new constraints to the models.




**Fig. 5** Preliminary angular distributions of the polarization observable $F$ from FROST in the $\gamma p \to K^+\Sigma^0$ reaction for various $W$ bins. Model curves are from the RPR-Ghent [29] (red), KAON-MAID [30] (blue), and Bonn-Gatchina [26] (magenta) models. Figure from N. Walford (U. Basel).

Many nucleon resonances in the mass region above 1.6 GeV decay predominantly through either $\pi\Delta$ or $\rho N$ intermediate states into $\pi\pi N$ final states. Double-pion photoproduction allows the study of resonances that have no significant coupling to the $\pi N$ channel with great potential to observe previously unobserved states. Many polarization observables are accessible in two-pion photoproduction off the nucleon [4]. The CLAS Collaboration was first to study the beam-helicity asymmetry $I^{\odot}$ for the two-pion-photoproduction reaction. The measurements covered energies between $W = 1.35$ and 2.30 GeV [31]. The FROST group is working on the determination of twelve different polarization observables from $\gamma p \to p\pi^+\pi^-$ data. As example, preliminary results are shown in Fig. 6 for observables that are accessible in measurements with unpolarized ($P_x$ and $P_y$) and circularly-polarized ($P_x^{\odot}$ and

$P_y^{\odot}$) photons off transversally-polarized protons. The data will strongly constrain coupled-channel analyses.




**Fig. 6** Preliminary CLAS results for the polarization observables $P_x$, $P_y$, $P_x^{\odot}$, and $P_y^{\odot}$ in the $\gamma p \to p\pi^+\pi^-$ reaction for 1.65 GeV $< W <$ 1.75 GeV. The angle $\phi*$ is the $\pi^+$ azimuthal angle in the rest frame of the $\pi^+\pi^-$ system with the $z$ direction along the total momentum of the $\pi^+\pi^-$ system. The data are fitted with even and odd low-order Fourier series according to the symmetry properties of the observables (red curves). Figure from A. Net (University of South Carolina).

Data have predominantly been taken off proton targets, and the new data from the FROST program will expand this data base over a large range of energy with many observables for polarized proton reactions. In constrast, data off neutrons are extremely sparse. However, measurements with both proton and neutron targets are needed to completely specify the amplitude of the reaction.

The CLAS collaboration has taken production data with circularly- and linearly-polarized photons off a polarized solid deuterium-hydride target (HDice) [8; 9] up to center-of-mass energies of $W \approx$ 2.3 GeV. The run conditions were optimized for polarized neutron reactions. The ongoing analyses of this run include single- and double-pion photoproduction and hyperon photoproduction off the bound neutron.

## 4 Conclusion

During the past years, knowledge of the baryon spectrum has increased greatly, as reflected in recent $N$ and $\Delta$ resonance updates in the Review of Particle Physics [32]. New polarized photoproduction data from CLAS off polarized and unpolarized proton and neutron targets are both under analysis and becoming available. These data will contribute to complete or nearly-complete experiments, and will challenge previously poorly constrained models. It is very likely that these data will have a tremendous impact on the understanding of baryon resonances and may provide evidence for new states found in coupled-channel analyses.

The quasi-real photon tagger for CLAS12 will allow to expand these photoproduction studies to higher beam energies with new experiments after the energy upgrade of JLab [33].

## References


1. V. Credé and W. Roberts, Progress towards understanding baryon resonances, Rept. Prog. Phys. **76** 076301 (2013).
2. R. G. Edwards, J. J. Dudek, D. G. Richards, and S. J. Wallace, Excited state baryon spectroscopy from lattice QCD, Phys. Rev. **D84** 074508 (2011).
3. W.-T. Chiang and F. Tabakin, Completeness rules for spin observables in pseudoscalar meson photoproduction, Phys. Rev. **C55** 2054 (1997).
4. W. Roberts and T. Oed, Polarization observables for two-pion production off the nucleon, Phys.Rev. **C71** 055201 (2005).
5. S. Strauch, Baryon spectroscopy with polarization observables from CLAS, in *12th Conference on the Intersections of Particle and Nuclear Physics (CIPANP 2015) Vail, Colorado, USA, May 19-24, 2015* (2015).
6. D. Sober *et al.*, The bremsstrahlung tagged photon beam in Hall B at JLab, Nucl. Instrum. Meth. A **440** 263 (2000).
7. C. Keith, J. Brock, C. Carlin, S. Comer, D. Kashy, J. McAndrew, D. Meekins, E. Pasyuk, J. Pierce, and M. Seely, The Jefferson Lab frozen spin target, Nucl. Instrum. Meth. A **684** 27 (2012).
8. A. M. Sandorfi, Unravelling the excitation spectrum of the nucleon, J. Phys. Conf. Ser. **424** 012001 (2013).
9. Jefferson Lab Experiment E06-101, "N* Resonances in Pseudoscalar-meson photo-production from Polarized Neutrons in $\mathbf{H} \cdot \mathbf{D}$ and a complete determination of the $\gamma n \to K^0 \Lambda$ amplitude", F. Klein and A.M. Sandorfi, spkespersons.
10. B. A. Mecking *et al.*, The CEBAF Large Acceptance Spectrometer (CLAS), Nucl. Instrum. Meth. **A503** 513 (2003).
11. M. Dugger *et al.*, Beam asymmetry for $\pi^+$ and $\pi^0$ photoproduction on the proton for photon energies from 1.102 to 1.862 GeV, Phys. Rev. C **88** 065203 (2013).
12. R. L. Workman, M. W. Paris, W. J. Briscoe, and I. I. Strakovsky, Unified Chew-Mandelstam SAID analysis of pion photoproduction data, Phys. Rev. **C86** 015202 (2012).
13. J. W. C. McNabb *et al.*, Hyperon photoproduction in the nucleon resonance region, Phys. Rev. **C69** 042201 (2004).
14. M. E. McCracken *et al.*, Differential cross section and recoil polarization measurements for the $\gamma p \to K^+ \Lambda$ reaction using CLAS at Jefferson Lab, Phys. Rev. **C81** 025201 (2010).
15. R. K. Bradford *et al.*, First measurement of beam-recoil observables $C_x$ and $C_z$ in hyperon photoproduction, Phys. Rev. **C75** 035205 (2007).
16. A. V. Sarantsev, V. A. Nikonov, A. V. Anisovich, E. Klempt, and U. Thoma, Decays of baryon resonances into $\Lambda K^+$, $\Sigma^0 K^+$ and $\Sigma^+ K^0$, Eur. Phys. J. **A25** 441 (2005).
17. V. A. Nikonov, A. V. Anisovich, E. Klempt, A. V. Sarantsev, and U. Thoma, Further evidence for $N(1900)P_{13}$ from photoproduction of hyperons, Phys. Lett. **B662** 245 (2008).
18. Jefferson Lab Experiment E02-112, "Search for Missing Nucleon Resonances in Hyperon Photoproduction", P. Eugenio, F. Klein, and L. Todor, spokespersons.
19. Jefferson Lab Experiment E03-105, "Pion Photoproduction from a Polarized Target", N. Benmouna, W. Briscoe, G. O'Rielly, I. Strakovsky, S. Strauch, spokespersons.
20. Jefferson Lab experiment E04-102, "Helicity Structure of Pion Photoproduction", D. Crabb, M. Khandaker, and D. Sober, spokespersons.
21. Jefferson Lab experiment E05-012, "Measurement of polarization observables in $\eta$-photoproduction with CLAS", M. Dugger and E. Pasyuk, spokespersons.
22. Jefferson Lab Experiment E06-013, "Measurement of $\pi^+\pi^-$ Photoproduction in Double-Polarization Experiments using CLAS", M. Bellis, V. Credé, S. Strauch, spokespersons.
23. S. Strauch *et al.*, First Measurement of the Polarization Observable $E$ in the $\mathbf{p}(\boldsymbol{\gamma}, \pi^+)n$ Reaction up to 2.25 GeV, Phys. Lett. **B750** 53 (2015).
24. A. V. Anisovich, V. Burkert, E. Klempt, V. A. Nikonov, E. Pasyuk, A. V. Sarantsev, S. Strauch, and U. Thoma, Existence of $\Delta(2200)7/2^-$ precludes chiral symmetry restoration at high mass, arXiv:1503.05774 [nucl-ex].

25. D. Rönchen, M. Döring, F. Huang, H. Haberzettl, J. Haidenbauer, C. Hanhart, S. Krewald, U. G. Meißner, and K. Nakayama, Photocouplings at the Pole from Pion Photoproduction, Eur. Phys. J. **A50** 101 (2014).
26. A. V. Anisovich, R. Beck, E. Klempt, V. A. Nikonov, A. V. Sarantsev, and U. Thoma, Properties of baryon resonances from a multichannel partial wave analysis, Eur. Phys. J. **A48** 15 (2012).
27. I. Senderovich *et al.*, First measurement of the helicity asymmetry $E$ in $\eta$ photoproduction on the proton, Phys. Lett. **B755** 64 (2016).
28. D. Drechsel, S. Kamalov, and L. Tiator, Unitary Isobar Model - MAID2007, Eur. Phys. J. **A34** 69 (2007).
29. T. Corthals, J. Ryckebusch, and T. Van Cauteren, Phys. Rev. **C73** (2006) 045207 and T. Vrancx, personal communication.
30. F. X. Lee, T. Mart, C. Bennhold, H. Haberzettl, and L. E. Wright, Nucl. Phys. **A695**, 237 (2001), "Kaon-MAID". Available from http://www.kph.uni-mainz.de/MAID/kaon/kaonmaid.html.
31. S. Strauch *et al.*, Beam-helicity asymmetries in double-charged-pion photoproduction on the proton, Phys. Rev. Lett. **95** 162003 (2005).
32. K. A. Olive *et al.*, Review of Particle Physics, Chin. Phys. **C38** 090001 (2014).
33. D. I. Glazier, A quasi-real photon tagger for CLAS12, Nucl. Phys. Proc. Suppl. **207-208** 204 (2010).